


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C05F 11/08, C02F 3/00, C05G 3/00		A1	(11) International Publication Number: WO 95/03259
			(43) International Publication Date: 2 February 1995 (02.02.95)
(21) International Application Number: PCT/US94/08431 (22) International Filing Date: 25 July 1994 (25.07.94) (30) Priority Data: 096,735 23 July 1993 (23.07.93) US (71) Applicant: IDAHO RESEARCH FOUNDATION, INC. [US/US]; 121 Sweet Avenue, Moscow, ID 83843-0178 (US). (72) Inventors: CRAWFORD, Donald, L.; 1988 Appaloosa Road, Moscow, ID 83843 (US). STEVENS, Todd, O.; Apartment 115, 50 Jadwin Avenue, Richland, WA 99352 (US). CRAWFORD, Ronald, L.; 4375 Broenneke Road, Moscow, ID 83843 (US). (74) Agents: STEPHENS, Donald, L., Jr. et al.; Klarquist, Sparkman, Campbell, Leigh & Whinston, One World Trade Center, Suite 1600, 121 S.W. Salmon Street, Portland, OR 97204 (US).		(81) Designated States: AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, HU, JP, KP, KR, KZ, LK, LU, LV, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: BIOLOGICAL SYSTEM FOR DEGRADING NITROAROMATICS IN WATER AND SOILS



(57) Abstract

Novel methods for biodegrading nitroaromatic compounds present as contaminants in soil or water using microorganisms are disclosed. Water is treatable directly; dry soil (22) is first converted into a fluid medium by addition of water. The preferred method comprises two stages, each employing microorganisms: a fermentative stage, followed by an anaerobic stage. The fermentative stage is rapid, wherein an inoculum of aerobic and/or facultative microorganisms (36) ferments a carbohydrate (30) added to the fluid medium, exhausting the oxygen in the fluid medium and thereby inhibiting oxidative polymerization of amino by-products of the nitroaromatics. In the subsequent anaerobic stage, an inoculum of a mixed population of anaerobic microorganisms (44) completes the mineralization of the contaminant nitroaromatics, using the remaining carbohydrate as a carbon and energy source. Preferably, the carbohydrate (30) is a starch and the aerobic and/or facultative microorganisms are amylolytic (36), which cleave the starch at a moderate rate throughout both stages, ensuring a sustained supply of metabolizable carbohydrates. The microorganisms are preferably selected to be resistant to the types and concentrations of nitroaromatics present as contaminants.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

BIOLOGICAL SYSTEM FOR DEGRADING NITROAROMATICS IN WATER AND SOILS

Field of the Invention

This invention relates to the biodegradation of various nitroaromatic compounds in
5 water and soils, including dinoseb (2-(1-methylpropyl)-4,6-dinitrophenol), using microorganisms.

Background of the Invention

Certain nitrophenolic compounds are sufficiently toxic to life to render them effective
for use as herbicides, insecticides, or miticides. Such compounds include dinoseb (2-(1-
methylpropyl)-4,6-dinitrophenol) which has been widely used as a herbicide since the 1950's on a
10 variety of crops in the United States. Concerns for the safety of agricultural workers has resulted
in discontinued use of dinoseb. However, numerous sites remain contaminated with this
compound.

Other nitroaromatic compounds are similar to dinoseb in terms of chemical structure,
but have other applications, such as in explosives. Such compounds include trinitrotoluene (TNT)
15 and dinitrotoluene (DNT). Because of the widespread use of these compounds over a lengthy
period of time, many sites have become contaminated with these compounds, including both
manufacturing and military sites.

Many nitroaromatic compounds are either poorly degradable or nondegradable in
field environments outside the laboratory. Previously, land farming was the favored method for
20 disposing of these and other chemicals, wherein the chemicals were mixed with soil, fertilizer
was added, and the mixture aerated to promote microbial activity. Unfortunately, nitroaromatics
were not satisfactorily degraded by land farming or other well-aerated processes. Possible
reasons include lack of nitroaromatic-degrading microorganisms, partitioning of the contaminant
chemicals to biologically sequestered or inhospitable parts of the environment, and accumulation
25 of toxic partial-breakdown by-products. Problems with land farming in general included the slow
rate of biodegradation, high expense, and accumulation of toxic by-products.

Other methods have been used to remove nitroaromatics and similar compounds from
contaminated soils, but with little practical success. Such methods include transportation of
contaminated soil to hazardous waste dumps, and on-site incineration of the soil. Problems with
30 such methods include high cost and poor accountability of the responsible party.

Previous laboratory studies indicated that certain nitroaromatic molecules are
susceptible at least to microbiological transformation. However, the studies did not disclose
biochemical mechanisms of such transformation or degradation or whether the nitroaromatic
compounds were completely mineralized. In one study, for example, a soil Moraxella
35 microorganism was isolated that was capable of growth on p-nitrophenol as its only source of
carbon and energy. Spain et al., Biochem. Biophys. Res. Comm. 88:634-641 (1959). In another
study, the anaerobic bacterium Veillonella alkalescens reductively transformed nitroaromatic

compounds, converting the nitro groups to amino groups. McCormack et al., Appl. Environ. Microbiol. **31**:949-958 (1976).

Aminoaromatic derivatives of nitroaromatics can undergo enzymatic oxidation to form polymeric (large molecular weight) materials. Parris, Residue Revs. **76**:1-30 (1980). In the field, such polymers are usually incorporated into soil humic matter. Channon et al., Biochem. J. **38**:70-85 (1944); McCormick et al., Appl. Environ. Microbiol. **31**:949-958 (1976); Simmons et al., Environ. Sci. Technol. **23**:115-121 (1989). Humic matter tends to be long-lived in soils, thereby representing a major long-term environmental fate of many nitroaromatics and aminoaromatics. Other soil microorganisms are capable of cleaving the azo linkages of polymerized aminoaromatics, often forming toxic by-products.

Bacteria are also able to attack nitrobenzoic acid, Cartwright and Cain, Biochem. J. **71**:248-261 (1959), as well as *o*-nitrophenol and *m*-nitrophenol, Zeyer and Kearney, J. Agric. Food Chem. **32**:238-242 (1984), where the nitro group is released as nitrite. Again, however, complete mineralization has not been demonstrated. Further, nitrite release has not been found to be a significant pathway for highly substituted nitroaromatics. No instance is currently known where a compound possessing more than one nitro substituent has been completely mineralized. In fact, the pertinent literature presents no evidence supporting ring cleavage of highly substituted nitroaromatics. Kaplan, "Biotransformation Pathways of Hazardous Energetic Organo-Nitro Compounds," in Biotechnology and Degradation. Adv. Appl. Biotechnol. Ser. 4:155-181, Gulf Pub. Co., Houston, TX (1990).

Aromatic groups in general appear to be degradable via only a few aerobic and anaerobic pathways. Gottschalk, Bacterial Metabolism, 2d ed., Springer Verlag, New York (1986), pp. 157-162; Berry et al., Microbiol. Rev. **51**:43-59 (1987); Schink, "Principles and Limits of Anaerobic Degradation: Environmental and Technological Aspects," in Zinder (ed.), Biology of Anaerobic Microorganisms, Wiley, New York (1988). Aerobically, many aromatic groups are degraded to catechol, protocatechuate or homogentisate by the action of oxygenase and dioxygenase enzymes. Catechol and protocatechuate can be degraded further by aromatic ring cleavage either *ortho* or *meta* to the hydroxyl groups. Because of the difficulty of working with anaerobic microorganisms and processes, biochemical pathways describing anaerobic degradation of aromatic compounds have been less well characterized.

Alkyl groups on aromatic rings are degradable via reactions similar to those for simple alkanes. Under aerobic conditions, the terminal carbon is oxidized to yield a carboxylic acid. Degradation then proceeds by β -cleavage to yield either benzoates (odd-numbered carbon chains) or phenylacetates (even-numbered carbon chains). No anaerobic microorganisms capable of carrying out this process have been isolated to date. In spite of the above results known in the art, there is little information currently available on practical means of using microbial cultures to bioremediate nitroaromatic-contaminated soils.

Dinoseb, an intensely yellow-colored compound visible at concentrations as low as 10 ppm, has been found to not significantly accumulate in agricultural soil at normal application rates, even after years of repeated application. Doyle et al., J. Agric. Food Chem. **26**:987-989 (1978). However, higher application rates, such as from spills of substantial amounts of the compound, can result in appreciable accumulation at a site. Presumably, therefore, dinoseb at lower concentrations is transformed by certain soil microorganisms. Such transformation appears to result only in the formation of amino and acetoamido forms of dinoseb, which apparently retain significant toxicity. Parris, Residue Revs. **76**:1-30 (1980).

Previous work on the biotransformation of the explosive 2,4,6-trinitrotoluene (TNT) indicates that the primary mode involves transformation (reduction) of the nitro groups. Kaplan, supra. A recent paper from Soviet researchers describes degradation of TNT by a strain of Pseudomonas fluorescens. Naumova et al., Mikrobiologiya **57**:218 (1988). But, while these reports shed some light on microbial events and hypothetical biochemical mechanisms therefor, they neither disclose nor suggest effective methods for bioremediating soils or wastewater contaminated with these compounds. Further, the Soviet results have not been confirmed outside the U.S.S.R.

Hence, although several anaerobic microbiological systems have been described for degrading other aromatic chemicals, little to no information is available on practical means of using these cultures to bioremediate contaminated soils and waters, especially soils and waters contaminated with nitroaromatics. In today's world, effective remediation of environmental sites contaminated with compounds, such as nitroaromatics, requires that the contaminants be completely mineralized to ensure the absence of latently toxic by-products. Such results for nitroaromatics simply have not been shown in the prior art, particularly as applicable to large-scale, low-cost bioremediation efforts.

Therefore, there remains a need for a method to effectively bioremediate dinoseb-contaminated soils, as well as soils contaminated with other nitroaromatic compounds, such as TNT and DNT.

Further, there is a need for such a method that can be performed at a natural site contaminated with dinoseb or a related nitroaromatic compound.

Further, there is a need for such a method that can completely degrade dinoseb and other nitroaromatics, leaving no detectable or environmentally significant amounts of aromatic by-products or other toxic intermediary compounds, including polymeric derivatives.

Further, there is a need for such a method employing microorganisms of types and species profiles normally found in many soil environments.

Further, there is a need for such a method that is inexpensive and easy to perform, particularly on a large-scale, in the field.

Further, there is a need for such a method that can be performed rapidly, including in the field.

Further, there is a need for such a method that will effect bioremediation of nitroaromatic-contaminated soil without specialized bioreactors or other complex equipment.

Summary of the Invention

5 In accordance with the present invention, soil or water contaminated with one or more nitroaromatic compounds is subjected to a two-stage bioremediation process employing different microorganisms during each stage. The stages comprise an initial fermentation stage followed by an anaerobic stage. Most of the actual biodegradation of the contaminant nitroaromatics takes place in the anaerobic stage. At the end of the anaerobic stage, the contaminant nitroaromatics have been biodegraded to non-toxic end-products.

10 As another aspect of the invention, complete biodegradation of nitroaromatics has been found to occur only under anaerobic conditions. Since anaerobiosis generally requires an aqueous environment, it is usually necessary to add extraneous water to a nitroaromatic-contaminated soil to create a fluid mud slurry of the soil before beginning the process. Contaminated water can be subjected to the process directly.

15 During the first stage of the process, the normally aerobic contaminated soil (with water added to form a mud slurry) or contaminated water alone is rendered anaerobic. The preferred method for achieving an anaerobic condition is by a fermentation of a supply of starchy carbohydrate or other readily fermentable carbon source added to the slurry or water. Fermentation, where the carbon source is a starchy carbohydrate, is performed by one or more
20 species of aerobic or facultatively anaerobic amylolytic microorganisms inoculated into the slurry or water. Amylolytic microorganisms are not required if the carbon source is a simple sugar, such as fructose or glucose.

As another aspect of the invention, the aerobic or facultative microorganisms are preferably isolated and enriched in a culture containing the particular nitroaromatic present in the
25 contaminated soil or water.

As another aspect of the invention, it is usually necessary at the beginning of the aerobic stage to add an extraneous source of nitrogen for the microorganisms. The nitrogen source is preferably in the form of ammonium ion or simple amino compounds readily utilizable by aerobic and anaerobic microorganisms.

30 As another aspect of the invention, it is preferable to stimulate a rapid onset of intense fermentation to quickly cause exhaustion of the oxygen dissolved in the slurry or water, thereby rendering the slurry or water anaerobic, without exhausting the carbon source. Quick attainment of anaerobiosis minimizes oxidative polymerization of any amino derivatives of the nitroaromatics that tend to form under aerobic conditions. Once formed, such polymers are
35 difficult to biodegrade. Rapid anaerobiosis can be achieved by inoculating the slurry or water with a large dosage of aerobic and/or facultative fermentative microorganisms.

If necessary, mineral nutrients, including phosphate salts, can be added to the soil slurry or water to facilitate microbial growth. Supplementary vitamins and cofactors may also be

required, but probably only when treating wastewaters having very little dissolved organic carbon. Soils generally have sufficient nutrients.

As another aspect of the invention, the carbon source added to the soil slurry or water for aerobic fermentation is preferably a starchy carbohydrate substance hydrolyzable to constituent sugars by amylolytic microorganisms in the aerobic inoculum. A starch is preferred over merely adding free sugar because the starch serves as a reservoir of metabolizable carbohydrate that ensures an adequate supply of easily metabolizable sugar for the microorganisms, both in the aerobic stage and in the anaerobic stage. Anaerobic biodegradation of nitroaromatics requires such a sustained sugar supply, however, too high sugar concentrations inhibit dinoseb degradation. A supply of sugar added at the beginning of the aerobic stage would generally be exhausted prematurely, making it difficult to maintain anaerobic conditions for the requisite amount of time to achieve complete mineralization of the nitroaromatics.

As another aspect of the invention, the amount of starchy carbohydrate to be added to the volume of soil slurry or water to be treated can be "tailored" to ensure the desired degree of biodegradation is attained and waste of carbohydrate is avoided. The amount of carbohydrate should be just sufficient to supply the metabolic needs of the microorganisms until the anaerobic biodegradation of contaminant nitroaromatics is complete.

As yet another aspect of the invention, once strict anaerobic conditions have been attained in the volume of slurry or water, an inoculum comprised of an anaerobic consortium of microorganisms is added to the volume to start the second, or anaerobic, stage. Anaerobic conditions are preferably determined via a potentiometric measurement, where a redox potential of -200 mV or less indicates strict anaerobic conditions.

As yet another aspect of the invention, the anaerobic consortium comprises multiple species of microorganisms that have been grown in a medium containing one or more nitroaromatics identical or similar to the contaminant nitroaromatics to be biodegraded. The anaerobic microorganisms are able to biodegrade the nitroaromatics in the presence of metabolizable sugar to simple non-toxic compounds, such as methane, carbon dioxide, and acetate.

As yet another aspect of the invention, after inoculation, the anaerobic consortium is afforded sufficient time to biodegrade the contaminant nitroaromatics in the soil slurry or water to non-toxic end-products. Because degradation of the nitroaromatic compounds occurs in an anaerobic environment, polymerization to toxic humic-like compounds and other large, long-lived, latently toxic molecules normally formed in aerobic environments is prevented.

As yet another aspect of the invention, the present method is preferably performed in a suitably large covered vessel for containing the contaminated soil slurry or water during bioremediation. Such containment ensures that anaerobic conditions in the soil slurry or water are reached more rapidly and are better controlled and maintained. Containment also facilitates

easier control of environmental parameters, such as temperature, pH, and, if needed, escape of volatile gases from the slurry or water being treated.

It is accordingly one object of the present invention to provide a method for effectively bioremediating soils and waters contaminated with one or more nitroaromatic compounds.

Another object of the present invention is to provide such a method that can be performed at natural sites contaminated with nitroaromatics.

Another object of the present invention is to provide such a method that will allow contaminant nitroaromatics in soil or water to be biodegraded to such an extent that no detectable or environmentally significant amounts of aromatic by-products or other toxic intermediary compounds are left in the soil or water, including latently toxic polymeric derivatives of the nitroaromatics.

Another object of the present invention is to provide such a method that utilizes microorganisms similar to those found in many soil and aquatic environments.

Another object is to provide such a method that is inexpensive and easy to perform, particularly on a large scale in the field.

Another object is to provide such a method that can be performed rapidly, even in the field.

These and other objects, features, and advantages of the present invention will become apparent with reference to the following description and drawing.

Brief Description of the Drawing

The drawing consists of multiple figures, in which:

FIG. 1 is a schematic depiction of the formation of an amino derivative of the representative nitroaromatic compound "dinoseb" under aerobic conditions and the subsequent aerobic polymerization of amino derivatives into undesirable humic-like compounds retaining the latent toxicity of dinoseb.

FIG. 2 is a graph showing the rate of biodegradation of dinoseb according to the present invention as a function of temperature.

FIG. 3 is a graph showing the relationship to dinoseb biodegradation rate of various concentrations of sugar (carbon), and ammonium chloride (nitrogen) for the anaerobic consortium of microorganisms.

FIG. 4 is a graph showing the rate of biodegradation of dinoseb according to the present invention as a function of pH.

FIG. 5 is a graph showing the kinetics of anaerobic biodegradation of dinoseb by the anaerobic consortium to intermediary compounds and the subsequent disappearance of the intermediary compounds.

FIG. 6 is a graph similar to FIG. 5 showing the kinetics of anaerobic biodegradation of 4,6-dinitro-*o*-cresol by an anaerobic consortium previously acclimated to dinoseb.

FIG. 7 is a graph similar to FIG. 5 showing the kinetics of anaerobic biodegradation of 3,5-dinitrobenzoate by an anaerobic consortium previously acclimated to dinoseb.

FIG. 8 is a graph similar to FIG. 5 showing the kinetics of anaerobic biodegradation of 2,4-dinitrotoluene by an anaerobic consortium previously acclimated to dinoseb.

5 FIG. 9 is a graph similar to FIG. 5 showing the kinetics of anaerobic biodegradation of 2,6-dinitrotoluene by an anaerobic consortium previously acclimated to dinoseb.

FIG. 10 is a graph similar to FIG. 5 showing the kinetics of anaerobic biodegradation of dinitrophenol by an anaerobic consortium previously acclimated to dinoseb.

10 FIG. 11 is a schematic depiction of a reaction pathway for the anaerobic biodegradation of dinoseb according to the process of the present invention.

FIG. 12 is a graph showing the effect of BESA, an inhibitor of methanogenesis, on the anaerobic biodegradation of dinoseb by an anaerobic consortium.

FIG. 13 is another schematic depiction of a reaction pathway for the anaerobic biodegradation of dinoseb according to the process of the present invention.

15 FIG. 14 is a representative schematic depiction of a process according to the present invention as it would be conducted at a field site.

Detailed Description of the Preferred Embodiment

I. Selection and Isolation of Dinoseb-Degrading Microorganisms

20 Various microorganisms capable of at least transforming dinoseb were preliminarily selected and enriched using a chemostat having a 1-L capacity vessel and agitation capability (Series 500 Fermenter, L. H. Fermentation, Hayward, CA). An approximately 250 mL volume of 3 mm diameter glass beads were placed in the bottom of the chemostat vessel to act as a soil-holding matrix. As a result, both aerated (supernatant liquid) and non-aerated (soil sediment) enrichment conditions were simultaneously maintained in the chemostat.

25 As used herein, "transformation" of dinoseb or other nitroaromatic compound means a chemical change other than degradation. The simplest way to confirm transformation of dinoseb is to observe the disappearance of the bright yellow color of the compound. Under aerobic conditions, dinoseb is transformed into an amino form that subsequently becomes polymerized by oxidative coupling. The term "degradation" denotes the complete mineralization of the subject nitroaromatic to methane, carbon dioxide, and acetate.

30 To provide a source of mineral nutrients to the microorganisms in the chemostat, a mineral nutrient solution was prepared containing the following solutes: KH_2PO_4 (272 mg/L), K_2HPO_4 (348 mg/L), Na_2SO_4 (5 mg/L), $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ (5 mg/L), $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (1 mg/L), and FeSO_4 (0.5 mg/L). The nutrient solution was supplemented with selected carbon and nitrogen sources, as discussed further below, in an attempt to effect a satisfactory preliminary selection of dinoseb-degrading microorganisms. The initial inoculum of microorganisms consisted of indigenous microbes associated with a 200-gram sample of a soil mixture removed from a site

previously exposed to dinoseb and suspected of having dinoseb-degrading activity. The chemostat was operated at a flow rate of 10 mL/hr, pH 7, and 25°C. Carbon and nitrogen sources for the microorganisms used individually in separate attempts at selection included: 50 ppm dinoseb plus 50 ppm 2,4-dinitrophenol, 50 ppm dinoseb plus 50 ppm 2,4-dinitrophenol plus 1 g/L NH_4Cl , 50 ppm dinoseb plus 50 ppm phenol, and 100 ppm dinoseb plus 0.5 g/L glucose and 1 g/L NH_4Cl .

No dinoseb degradation or turbidity occurred in the chemostat during thirty days' incubation and agitation with 50 ppm dinoseb plus 50 ppm 2,4-dinitrophenol as the sole carbon substrate. When the chemostat was operated for another thirty days with 50 ppm dinoseb plus 50 ppm 2,4-dinitrophenol and 1 g/L NH_4Cl , no dinoseb degradation or turbidity resulted. When the chemostat was operated for forty-two days with 50 ppm dinoseb, 50 ppm 2,4-dinitrophenol, and 50 ppm phenol, no dinoseb degradation occurred, but turbidity did develop. Most importantly, when the chemostat was operated with 0.5 g/L glucose, 100 ppm dinoseb, and 1 g/L NH_4Cl , turbidity developed immediately and dinoseb degradation began after twenty days. The latter result indicated that, to effect degradation of dinoseb by either aerobic or anaerobic processes, supplementary glucose is required as a carbon source. Continuing with the latter operational conditions, the flow rate was then increased to 20 mL/hr for another thirty days to complete the preliminary selection process.

After preliminary selection, samples of the supernatant liquid presumed to contain various aerobic and/or facultative microorganisms were removed from the chemostat and plated on nutrient agar to obtain a number of isolates of specific microorganisms. The isolates were then cultured under aerobic, anaerobic, and microaerophilic conditions wherein their individual abilities to transform or degrade dinoseb were evaluated. For aerobic culturing, the mineral nutrient solution described above was used, supplemented with the following: dinoseb (50 mg/L), glucose or fructose (500 mg/L), NH_4Cl (1.0 g/L), $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (0.5 mg/L), H_3BO_3 (0.05 mg/L), ZnCl_2 (0.05 mg/L), CuCl_2 (0.03 mg/L), $\text{Na}_2\text{MoO}_4 \cdot 2\text{H}_2\text{O}$ (0.01 mg/L), $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ (0.5 mg/L), $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (0.05 mg/L), Na_2SeO_3 (0.05 mg/L), and a vitamin solution recommended by Wolin et al., *J. Biol. Chem.* 238:2882-2886 (1963). Yeast extract was added to the culture solution to a concentration of 0.5 g/L. Yeast extract served as a convenient source of additional carbon and energy for the microorganisms, as well as a source of additional vitamins and cofactors.

For culturing under denitrifying (anaerobic) conditions, the aerobic culture medium was supplemented with 1 g/L KNO_3 , boiled under nitrogen gas, and sealed in glass containers with butyl rubber stoppers before inoculation. Culturing under denitrifying conditions is a way of selecting for facultative anaerobes. Under such conditions, nitrate is employed by the respiring microbes as an electron acceptor rather than oxygen as used by respiring aerobic microorganisms. Facultative anaerobes can be cultured in either aerobic or anaerobic environments.

A reduced anaerobic mineral medium (RAMM) was used for culturing the microorganisms under anaerobic conditions. RAMM comprised the same ingredients as listed

above for aerobic cultures, but with 10 mg/L resazurin added as a redox indicator, 10 mg/L $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$ added as a reducing agent, and 1.2 g/L NaHCO_3 . (Preferably, 0.1 g/L yeast extract is also added.) Anaerobic cultures were grown in serum bottles and balch tubes sealed with butyl rubber stoppers, using strict anaerobic procedures, as detailed in Ljungdahl and

- 5 Wiegel, "Working with Anaerobic Bacteria," in Demain and Solomon (eds.), Manual of Industrial Microbiology and Biotechnology, American Society for Microbiology, Washington, D.C. (1986) pp. 84-96.

Several isolates of microorganisms which could aerobically transform dinoseb were obtained from the supernatant liquid of the chemostat containing glucose, dinoseb, and NH_4Cl .

- 10 Characteristics of these isolates are presented in Table 1.

Table 1
Isolates That Aerobically Transform Dinoseb

15	Isolate	Gram Reaction	Shape	Colony ^a Morphol.	Oxidase	Catalase	Facultative	Dinoseb Transformation When: microaerophilic	denitrifying
	TDN-1	+	rods	R,O,P	+	+	no	yes	no
	TDN-2	-	rods	I,T	+/-	+	yes	yes	yes
20	TDN-3	-	rods	R,T	-	+	yes	yes ^b	no ^b
	TDN-4	+/-	rods	R,Y	+	+	yes	yes	no
25	TDN-5	+/-	cocci	S,T	+/-	+	yes	yes	yes

^a = Colonial Morphologies: R = round, I = irregular, S = spreading, T = transparent, O = opaque, Y = yellow, P = produces yellow-green pigment.

^b = Dinoseb transformation in this strain is inhibited by nitrate

30

- Under microaerophilic conditions, every strain caused the dinoseb-containing culture medium to turn bright red. Such transformation was not noted with any of the strains in aerated
- 35 dinoseb-containing cultures. As used herein, the term "microaerophilic conditions" refers to a culture environment having an appreciably depressed concentration of dissolved oxygen as compared to aerobic conditions, but not so low as to be strictly anaerobic. Under such conditions, aminoaromatics can still undergo polymerization reactions. This is in contrast to strictly anaerobic conditions under which such polymerization reactions are blocked.
- 40 Microorganisms termed "microaerophiles" undergo optimal growth under microaerophilic conditions.

- The red substance obtained under microaerophilic conditions with all the strains could not be extracted with organic solvents, nor could it be resolved by thin-layer chromatography (TLC). Analysis suggested that the red substance contained the amino
- 45 derivatives of dinoseb. After two to three weeks, the red color faded and a brown precipitate formed. TLC analysis of the brown precipitate showed a continuous smear with no discernable bands. These results indicate that dinoseb is transformed under microaerophilic conditions to an

amino form which is oxidatively polymerized to larger random-length molecules and not degraded.

Strains TDN-2 and TDN-5 appeared to be facultative anaerobes and were able to carry out the transformation of dinoseb to the red metabolite under denitrifying (anaerobic) conditions, in which case the brown precipitate did not form.

To quantify the transformation of dinoseb by the strains of Table 1, dinoseb concentrations were determined by High Performance Liquid Chromatography (HPLC) using a binary gradient of 10% tetrahydrofuran and methanol (solution A) and 1% acetic acid and water (solution B) on a 250 x 2 mm Phenomenex "Spherex" 5 μ m C18 reverse-phase column (Phenomenex Corp., Rancho Palos Verdes, CA). A Hewlett-Packard Model 1090A instrument, equipped with a diode-array detector and a computerized data system, was used for the analyses. The solvent flow rate was 0.4 mL/min, and the column temperature was 40°C. The gradient program was a ten-minute gradient from 60% solution A plus 40% solution B to 100% solution A, followed by five minutes at 100% solution A. Detection of dinoseb and transformation products thereof was by use of the diode-array detector, measuring UV absorption at 268, 225, and 385 nm, with continuous scanning of the absorption spectrum from 190 to 450 nm.

HPLC analysis of the culture medium from strains TDN-2 and TDN-5, of Table 1 also showed accumulation of a single dinoseb-transformation product with a HPLC retention time of 2.05 minutes. Dinoseb transformation by strain TDN-3 was inhibited by nitrate in both microaerophilic and denitrifying conditions. No dinoseb transformation occurred in anaerobic cultures of TDN-4, but transformation did occur when the flasks were opened and the cultures exposed to air. Strain TDN-1 was obligately aerobic and only transformed dinoseb under microaerophilic conditions.

As indicated in Table 1, these isolates of dinoseb-transforming microorganisms were taxonomically diverse, although no definite species identifications were made. Despite this apparent species diversity, the isolates appeared to carry out similar reactions when transforming dinoseb.

The above results indicate that, in oxygen-containing environments, the isolates obtained from the aerobic supernatant of the chemostat only reduced the nitro groups of dinoseb, thereby forming amino products that were subsequently oxidized by extracellular enzymes to form amorphous polymeric compounds, as diagrammed in FIG. 1. Since the dinoseb is apparently not actually degraded via such an aerobic process, but merely "transformed" into an amorphous polymer, the products of the process probably latently retain all the toxicity of dinoseb. Thus, it appears that aerobic bacteria such as certain of the strains listed in Table 1 do not actually detoxify dinoseb and would, therefore, not be appropriate for use in bioremediation of nitroaromatic contaminants in wastewaters or contaminated soils.

Although the above results were useful in elucidating the mechanism of dinoseb transformation in aerobic environments, it became clear that a biological method for degrading

dinoseb and related nitroaromatic compounds from contaminated soils and waters must include the use of microorganisms selected in a dinoseb-containing anaerobic environment. As a result, microorganisms resident in the anaerobic sediment of the chemostat supplied with medium containing 100 ppm dinoseb plus 0.5 g/L glucose and 1 g/L NH_4Cl were cultured and tested.

5 A consortium (stable mixed population) of anaerobic microorganisms capable of degrading dinoseb to non-aromatic products was enriched from the population of such organisms in the chemostat as follows. Sediment from the chemostat consisting of soil was used to inoculate strictly anaerobic medium comprised of the mineral nutrient solution described above with added 1 g/L fructose, 1 g/L NH_4Cl , and 100 ppm dinoseb. After five weeks' incubation, the bright
10 yellow color of the medium (due to the presence of dinoseb) changed to a bright orange, then faded to colorlessness followed by development of turbidity. This activity could be maintained in mineral medium for three or four sediment-free transfers, but not in medium containing 0.2 g/L yeast extract or 5% rumen fluid. Sediment-free anaerobic dinoseb-degrading cultures could be maintained indefinitely by making three transfers in mineral medium, followed by one transfer in
15 yeast extract-containing medium, followed by further mineral medium transfers. After eighteen months of such transfers, the dinoseb-degrading cultures remained stable in the yeast extract-containing medium, which resulted in 5- to 10-fold faster degradation of dinoseb than in mineral medium without yeast extract.

Degradation of dinoseb to non-aromatic products by the anaerobic consortium did
20 not occur unless strict anaerobic procedures were followed during preparation of the media and during culture transfers. This indicates that actual dinoseb degradation to non-aromatic products, as opposed to mere transformation, is an anaerobic process.

Several parameters should be controlled for optimal dinoseb degradation by the anaerobic consortium. These include temperature, nitrogen and pH.

25 As shown in FIG. 2, the optimal temperature for dinoseb degradation is about 25°C. Although FIG. 2 only covers a range from 20°C to 40°C, a useful temperature range would be within the range for mesophilic microorganisms, generally between 10°C and 40°C. Temperatures higher than about 40°C would either kill important microorganisms or shut down key enzymatic reactions. Lower temperatures within this range, including temperatures within the
30 range 10°C to 20°C, would merely result in a slower metabolic rate of the microorganisms, the rate generally dropping by about half for every ten-degree drop in temperature.

As shown in FIG. 3, the optimal sugar concentration (either fructose, glucose, or other simple sugar is suitable) is about 0.5 g/L, as indicated by the "carbon" line. The optimal NH_4Cl concentration is about 4 g/L, as indicated by the "nitrogen" line. However, the anaerobic
35 culture is largely insensitive to NH_4Cl concentration between about 1 and 5 g/L. NH_4Cl serves as an important source of nitrogen for the anaerobic microorganisms. Other nitrogen-containing compounds can also be used, so long as the nitrogen is in a form such as ammonium ion or

amino groups. The nitrogen source should not be a nitrate because nitrates may inhibit the process of nitroaromatic degradation by these microorganisms.

Finally, the optimal pH is about 7, as shown in FIG. 4, but the anaerobic culture appears to be largely insensitive to pH values between 6 and 8.

5 The anaerobic consortium degraded dinoseb via a series of intermediate aromatic products (A, B, C, and D, as shown in FIG. 5) which could be detected by HPLC. As determined by UV absorption spectra, no further aromatic products could be detected after thirty days, indicating that complete cleavage of the aromatic ring occurred. Such aromatic cleavage is a key step in the degradation of dinoseb to non-toxic compounds.

10 The stable consortium of anaerobic microorganisms contained at least three bacterial morphologies, including: short coccobacilli, 1-1.5 μm long; medium-sized rods, .75x2 μm ; and large rods, 1-1.5x4 μm . Briefly exposing the consortium to air before anaerobic incubation resulted in elimination of the large rods. Also, dinoseb degradation did not proceed beyond intermediate D (FIG. 5). When the anaerobic consortium was used to inoculate dinoseb-
15 containing aerobic media, a single bacterial species grew. That species was a gram-negative rod which grew as a coccobacillus under anaerobic conditions. The coccobacillus transformed dinoseb to a single uncharacterized product in aerobic cultures and to a different product in anaerobic cultures. The bacterium also was catalase positive, oxidase negative, and most closely matched *Klebsiella oxytoca* (similarity index 0.677), using the BioLog GN identification system
20 (Hayward, CA). The BioLog GN identification system is a method of testing bacteria for utilization of ninety-five different carbon substrates, where the resulting pattern growth is compared with a database of patterns for known species of bacteria.

 The anaerobic consortium was tested for its ability to degrade other nitroaromatic compounds. Utilization of other nitroaromatic substrates was determined by growing cultures in
25 medium similar to that described above, but with 50 ppm of the appropriate nitroaromatic compound substituted for dinoseb. The anaerobic consortium was able to completely degrade 4,6-dinitro-*o*-cresol (DNOC, FIG. 6) and 3,5-dinitrobenzoate (3,5-DNB, FIG. 7) to non-aromatic compounds. 2,4-Dinitrotoluene (2,4-DNT, FIG. 8) and 2,6-dinitrotoluene (2,6-DNT, FIG. 9)
30 were degraded to intermediate products, but it was unclear whether or not the products were eventually degraded to non-aromatics. After sixty days, the concentration of 2,4-dinitrophenol (2,4-DNP, FIG. 10) began to decline, but the parent compound persisted for at least four months in these cultures. Degradation of these other nitroaromatic compounds was much slower than for dinoseb.

 A dinoseb anaerobic degradation pathway consistent with the above results is shown
35 in FIG. 11.

 When bromoethanesulfonic acid (BESA), a specific inhibitor of methanogenesis, was added to the anaerobic consortium at 200 μM concentration, dinoseb degradation was slowed and products C and D accumulated (FIG. 12). These products remained in the culture medium

for at least another three months. Accumulation of C indicated that D is formed from C and that BESA at least partially blocked formation of D. Therefore, the reaction from C to D is probably a hydrogen-generating reaction. Since the reaction results in an increase in hydrophobicity, it probably involves removal of one or both acetamide groups.

5 The anaerobic dinoseb-degradation intermediates were tentatively identified as follows. Anaerobic dinoseb cultures were extracted and subjected to TLC, as described above, during both the early stages (orange color) and later stages (colorless) of dinoseb degradation. Extracts from the early stages yielded two TLC bands which were not present in uninoculated controls. GCMS analysis indicated that both bands contained multiple compounds. Similar
10 analysis of the extract from the later stages of degradation yielded one band not present in uninoculated controls. The mass spectra showed fragmentation patterns similar to that for dinoseb, confirming that they correspond to dinoseb derivatives. The intermediates as identified and/or hypothesized are shown in FIG. 11.

15 The molecular formulas of the intermediates, based on isotope abundance calculations from the molecular weights (± 0.005 AMU), are shown in Table 2. The intermediate with molecular weight 220 could not be assigned a molecular formula; however, the stated mass was expected from 2-sec-butyl-4-nitro-6-aminophenol. The major product in the later stages of degradation was identified as 2-aminobenzoic acid (anthranilic acid) by GCMS comparison with authentic standards.

-14-

Table 2

**Accurate Mass Determinations and Molecular
Formulas for Compounds Detected by GC/MS**

	<u>source</u>	<u>molecular mass</u>	<u>standard deviation</u>	<u>molecular formula</u>	<u>deviation from observed mass</u>
5	dinoseb	240.0699	.0084	C ₁₀ H ₁₂ N ₂ O ₅	.0047
	band 1	210.0891	.0063	NR	
10	band 1	220.0998	.0081	NR	
	band 1	264.0995	.0065	NR	
15	band 2	234.0966	.0071	C ₁₂ H ₁₄ N ₂ O ₃	.0038
	band 2	165.1147	.0046	C ₁₀ H ₁₃ NO	.0007
20	band 1B	136.060	.0088	C ₇ H ₈ N ₂ O	.0035
	band 1B	137.0508	.0047	C ₇ H ₇ NO ₂	
25	NR - no reasonable formula could be calculated from this mass reading				

Under anaerobic conditions the reduced dinoseb derivatives would not be subject to the oxidative coupling reactions found in the aerobic environment. Partially degraded molecules persisting as dissolved monomers are, therefore, available for further biodegradation by various anaerobic bacteria.

It is unclear why an external carbon source is required for dinoseb mineralization by the anaerobic consortium of microorganisms. One or more of these microorganisms may effect certain chemical changes to the dinoseb molecule, but derive no metabolic energy from such reactions or from the products of such reactions. Alternatively, microorganisms that carry out later steps in the degradation process might be inhibited by dinoseb. In the latter case, ring cleavage products would not be available for use by bacteria carrying out early steps in the degradation process.

The results described above indicate, therefore, that dinoseb is biodegradable by the anaerobic consortium of microorganisms. The initial step involving reduction of nitro groups to amino groups appears to occur in a manner similar to that for rumen microorganisms. Intermediates detected having molecular weights of 220, 234, and 264 were not identified, but probably represent N-alkylated forms of reduced dinoseb. The alkyl side chain of dinoseb is also anaerobically removed, as indicated by formation of anthranilic acid. FIG. 13 shows a dinoseb degradation pathway consistent with the results above.

Studies in which [¹⁴C]-ring-labeled dinoseb was degraded as above yielded [¹⁴C]-acetate, indicating that dinoseb rings are cleaved. Additionally, we have found that the presence of nitroaromatics inhibits methanogenesis. However, the nitro groups are readily reduced to amino groups, after which methanogenesis resumes. The only intermediates identified to date are

amino compounds, but numerous other "intermediates" have been found that remain to be specifically identified.

When tested with other nitroaromatic substrates, the anaerobic consortium was able to completely degrade nitrocresols or nitrobenzoates, but was unable to cleave the aromatic ring of nitrotoluenes or to degrade nitrophenols. It is probable that other selections performed in a manner similar to that described above would give rise to other anaerobic populations which can utilize these compounds. For example, passing a medium containing a nitrophenolic compound through a chemostat, as described above, inoculated with a diverse population of soil microorganisms would be expected to select for anaerobes capable of mineralizing nitrophenols. In other words, anaerobic consortia can be "tailored" for degrading a particular type of nitroaromatic compound.

II. Bioremediation of Fluid Media

As used herein, the term "fluid medium" refers to waters and slurries, including mud, comprising water plus soil or other particulate material.

The process of the present invention requires an aqueous fluid medium because anaerobiosis can only occur in aqueous environments. As a result, bioremediation of a contaminated dry soil requires that water be added, forming a mud slurry, to provide a sufficiently aqueous environment for metabolic activity by the anaerobic consortium of microorganisms. Such an aqueous environment is automatically provided when the process is to be applied to a contaminated wastewater.

In addition, the anaerobic consortium is unable to withstand prolonged exposure to oxygen. Consequently, an inoculum comprised of such microorganisms should not be added to a volume of soil slurry or wastewater to be subjected to biodegradation without first rendering the volume anaerobic. The transition to anaerobiosis should be as rapid as possible to preclude aerobic oxidative coupling of the contaminant nitroaromatics to polymeric forms via the action of indigenous aerobic microorganisms.

For the experiments described below, the anaerobic dinoseb-degrading consortium isolated as described above was maintained on a reduced anaerobic mineral medium (RAMM) consisting of: KH_2PO_4 (0.27 g/L), K_2HPO_4 (0.35 g/L), NH_4Cl (1.5 g/L), glucose (0.5 g/L), yeast extract (0.1 g/L), $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (15 mg/L), $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ (20 mg/L), $\text{FeCl}_2 \cdot 2\text{H}_2\text{O}$ (4 mg/L), $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$ (0.5 mg/L), H_3BO_3 (0.05 mg/L), ZnCl_2 (0.05 mg/L), $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ (0.05 mg/L), $\text{NiCl}_2 \cdot 6\text{H}_2\text{O}$ (0.05 mg/L), CuCl_2 (0.03 mg/L), $\text{NaMoO}_4 \cdot 2\text{H}_2\text{O}$ (0.01 mg/L), NaHCO_3 (2.4 g/L), and 1 ng/L resazurin. Cultures were incubated in darkness without shaking at 30°C. Strict anaerobic procedures were followed during all media preparations and transfer operations. Ljungdahl and Wiegel, "Working with Anaerobic Bacteria," in Demain and Solomon (eds.), Manual of Industrial Microbiology and Biotechnology, American Society for Microbiology, Washington, D.C. (1986).

In order to sustain anaerobiosis in the present process, a source of readily-metabolizable carbon, such as sugar, is required as a supplement. Preferably, the carbon source is a complex carbohydrate from which sugars are "released" over a period of time, rather than a supply of sugar that is completely "available" for immediate consumption. Because complex carbohydrates must be enzymatically cleaved to yield metabolizable sugar, they allow maintenance of strict anaerobic conditions in the fluid medium to be extended for a time period sufficient to biodegrade the particular concentration of contaminant nitroaromatic without the need to add more carbon source. For example, if a large amount of an "available" sugar, such as glucose or fructose, were added to an aerobic aqueous slurry of soil, anaerobiosis would be quickly achieved. However, maintenance of strict anaerobiosis for the several weeks that may be required to achieve complete degradation of the nitroaromatic would be difficult due to an initially very high rate of sugar metabolism followed by premature exhaustion of the sugar supply.

Although sugar could be added at one or more additional times during maintenance of anaerobic conditions before complete nitroaromatic degradation was achieved, the added sugar would have to be added in controlled amounts at specific times and mixed each time into the fluid being treated. Such adding and mixing impart unnecessary complexity to the method. Further, use of more complex carbohydrates, such as starch, as a carbon source results in a steadier degradation rate. Repeated additions of sugar result in an undulating rate which is less efficient.

In view of the above, desired characteristics of the carbon source include a high energy content, ready metabolizability, low numbers of indigenous heterotrophic bacteria which might compete with the anaerobic inoculum, and sustained metabolic availability sufficient to maintain prolonged steady rate of anaerobiosis. On the basis of these criteria, a hydrolyzable polysaccharide such as starch is a more suitable carbon source than free sugar.

After evaluation of a number of starchy by-products from various food processing plants, a preparation of dewatered solids from a potato processing plant was selected as the preferred carbon source. The characteristics of the potato waste included: 42% solids, 215 mg/g available starch, 6.7 mg/g total nitrogen, 2.6×10^4 culturable heterotrophic bacteria per gram, and 8×10^3 culturable amylolytic (able to hydrolyze starch) bacteria per gram. Chief advantages of starchy potato waste are available in large amounts and at a low cost. However, any vegetable or grain starch would probably suffice.

Candidate starchy carbohydrates were analyzed for solid content by weighing after oven drying, and for total nitrogen content by the Kjeldahl procedure commonly known in the art. Available starch was determined by incubating sterilized 1-gram samples with 300 units of α -amylase and 100 μ L of diazyme L-100 (Miles Pharmaceuticals, Elkhart, IN) for twenty-four hours in 10 mL of sterile 0.4 M phosphate buffer, pH 7.0. After incubation, the samples were diluted to 100 mL and analyzed for reducing sugars using the dinitrosalicylate assay with similarly prepared glucose as a standard. Miller, *Anal. Chem.* 31:426-428 (1959). Available starch was assayed as mg sugar released per gram dry weight of starchy carbohydrate. Numbers

and types of bacteria in starchy carbohydrates were determined by standard plate counts in aerobic mineral medium agar containing (for total heterotrophic counts) 2 g glucose, 0.4 g yeast extract, 1 g NH_4Cl , and 1 g NaNO_3 per liter, or (for amylolytic bacteria) 2 g soluble starch, 1 g NH_4Cl , and 1 g NaNO_3 per liter for amylolytic bacteria.

5 Since the anaerobic consortium cannot withstand prolonged exposure to air, it is necessary to pretreat the volume of wastewater or soil slurry with a rapid aerobic fermentation step to deplete oxygen in the volume and achieve anaerobiosis before adding the anaerobic consortium. The transition from aerobic fermentation to anaerobiosis should be as rapid as possible to preclude oxidative coupling of aminoaromatic derivatives of the contaminant
10 nitroaromatic in the aerobic step. According to the results of tests performed using loamy sand and rich silt-loam soils, simply flooding the soil with water does not produce sufficiently rapid anaerobiosis. Mere addition of sugar produces rapid anaerobiosis by an initial high rate of aerobic fermentation that rapidly consumes the available oxygen in the liquid. However, as discussed above, use of sugar as a carbon source instead of a complex carbohydrate, such as
15 starch, usually results in premature exhaustion of the carbon source. Also, even with elevated concentrations of added sugar, at least partially aerobic conditions often reform before completion of the nitroaromatic degradation. Use of dewatered potato solids or other starchy carbohydrate that is metabolized more slowly than free sugar results in a satisfactorily rapid anaerobiosis following aerobic fermentation, while yielding a sustained sugar concentration over the period of
20 time required to completely degrade the contaminant nitroaromatic.

 To evaluate the relative effectiveness of starchy carbohydrates as a carbon and energy source versus a sugar such as glucose alone, 300-g samples of various types of soil, such as loamy sand and rich silt-loam, were individually placed in one-liter Erlenmeyer flasks and flooded with 200 mL of 0.4 M phosphate buffer (pH 7) to form a mud slurry. Flasks were
25 covered with aluminum foil and incubated without shaking at 25°C. At various times during incubation, both pH and redox potential of the slurry were measured potentiometrically at 0, 1, and 3 cm beneath the liquid surface. A 1 mL sample was also removed for each analysis of residual sugar. A redox potential of -200 mV or less was indicative of a strictly anaerobic condition in the liquid.

30 The amount of starchy carbohydrate to add to a volume of wastewater or mud slurry for particulate dosages of fermentative and anaerobic organisms may have to be determined experimentally and optimized for a particular soil or wastewater and the particular type and concentration of contaminant nitroaromatic. To reduce costs in the actual bioremediation process, the amount of starch should be determined that will just sustain anaerobic conditions for the time
35 required to achieve the desired degree of nitroaromatic degradation and no longer.

 Any of a number of species of fermentative amylolytic microorganisms indigenously present in the starchy carbohydrate can serve to hydrolytically cleave the starch into constituent sugars when the starch is added to a volume of wastewater or slurry of soil. Such fermentative

microorganisms would include aerobic and/or facultative microorganisms. Starch cleavage occurs via amylase enzymes secreted by the microorganisms into the surrounding aqueous medium.

Although amylase action is optimal in aerobic environments, the enzyme will continue to hydrolyze starch when the medium becomes anaerobic. Additionally, many facultative amylolytic
5 microorganisms will survive and continue to secrete amylase after a medium has been converted from aerobic to anaerobic. If sugar is used as the carbon source, the fermentative microorganisms need not be amylolytic.

In a series of incubation tests where the soil contained 25 ppm dinoseb, anaerobiosis occurred only after several days in the presence of the starchy potato by-product, where the initial
10 aerobic fermentation was performed by fermentative amylolytic bacteria (representing a number of species) indigenous to the starchy potato by-product. Several days to achieve anaerobiosis is too long. Evidently, the indigenous bacterial population was adversely stressed by the dinoseb and unable to metabolically respond in a rapid manner. To selectively enrich for dinoseb-resistant amylolytic bacteria, additional starchy potato by-product and dinoseb (to 100 ppm) were
15 added to a flask from these experiments. After a prolonged incubation to achieve selection, several dinoseb-resistant bacteria were isolated from the flask. One strain (strain DSA-1), identified as facultative Klebsiella oxytoca by the BioLog GN system, retained good amylolytic activity in the presence of 100 ppm dinoseb.

When dinoseb-containing soil received an inoculum of Klebsiella oxytoca strain
20 DSA-1, along with the starchy potato by-product, a greatly reduced time to achieve anaerobiosis was noted. In the latter case, anaerobiosis was established almost as fast as in control soils lacking dinoseb. Klebsiella oxytoca strain DSA-1 was thus regarded as a preferred inoculation strain with which to achieve anaerobiosis for degrading dinoseb in the presence of a starchy carbohydrate.

Although Klebsiella oxytoca strain DSA-1 was the particular bacterium selected for
25 in the above-described experiments involving dinoseb, it is expected that other selections performed in a similar manner using a source of starch containing indigenous aerobic and/or facultative "fermentative" microorganisms will probably yield other species and strains after selection. Additionally, it is expected that using a nitroaromatic other than dinoseb for supplying
30 the selection pressure would probably give rise to other satisfactory fermentative species and/or strains resistant to the particular nitroaromatic. The lesson from these studies is that the indigenous microflora associated with starches are probably not sufficiently tolerant to most nitroaromatics to facilitate the required high initial rate of fermentation to achieve rapid anaerobiosis in a volume of soil slurry or wastewater containing nitroaromatic compounds. As a
35 result, it will probably be necessary to employ an aerobic or facultative species and strain preselected against the particular nitroaromatics to be biodegraded.

In order to achieve anaerobiosis in the shortest amount of time, it is preferable to inoculate using a large dose of preselected fermentative microorganisms. Our studies indicate that

a dose of about 10^7 to 10^8 CFU per gram dry soil or mL water is an effective dose. Cost constraints generally preclude larger doses. Also, larger doses generally do not achieve proportionately shorter times to anaerobiosis.

In another series of tests, bioremediations of 1-kg samples of each type of dinoseb-contaminated soil (loamy sand and silt-loam) were individually performed by adding water, starchy potato by-product, and pretreating the resulting slurry with a large inoculum (as above) of *Klebsiella oxytoca* strain DSA-1. After each slurry was rendered anaerobic (redox potential (-200 mV), it was inoculated with a similarly large dose of the dinoseb-degrading consortium of anaerobic microorganisms. Dinoseb was found to be completely converted to intermediate compounds within one week of anaerobic inoculation. After four weeks, the concentrations of the intermediate compounds declined below detectable limits. At the end of the four-week period, only 0.5 ppm of dinoseb could be Soxhlet-extracted from the treated loamy sand soil, and none could be extracted from the treated silt loam. By contrast, in the uninoculated loamy sand control, dinoseb did not decline. In the uninoculated silt loam control, dinoseb did decline after several days' lag time, but no intermediate compounds metabolically derived from dinoseb were detected.

In similar experiments, soils contaminated with like concentrations of 4,6-dinitro-*p*-cresol and 3,5-dinitrobenzoate were bioremediated within thirty days.

To perform the above experiments, soils were adjusted to 100 ppm dinoseb before adding any microorganisms or starchy carbohydrate by adding to the soil a solution of dinoseb in methanol and allowing the soil to dry completely afterward. For each soil sample, 1 kg dinoseb-supplemented soil was mixed with 2 g of the starchy carbohydrate. The mixture was inoculated with the amylolytic bacteria, placed in an open two-liter capacity Erlenmeyer flask, and a sufficient volume 0.4 M phosphate buffer (pH 7) was added to saturate the water-holding capacity of the soil. When the redox potential of the soil solution became less than -200 mV, a 50 mL volume (O.D. ≈ 1) of the anaerobic consortium was injected below the surface of the soil. Samples (approximately 5 g each) were removed periodically, weighed and vortexed with 5 mL 0.1 N NaOH. Samples were centrifuged to remove the solids, and the supernatant was analyzed for dinoseb and its metabolites. At the end of the incubation, the remaining soil was extracted and analyzed for dinoseb.

Dinoseb concentrations were determined by high performance liquid chromatography (HPLC) using a binary gradient of 10% tetrahydrofuran in methanol (solution A) and 1% acetic acid in water (solution B) on a 250x2 mm Phenomenex "Spherex" 5 μ m C18 reverse-phase column. Analyses were performed using a Hewlett-Packard Model 1090A instrument, as described above. Dinoseb was extracted from soils by Soxhlet extraction with ethyl acetate for five hours. Before extraction, each sample was amended with 200 μ L of a 0.25% solution of 4,6-dinitro-*p*-cresol in methanol, which served as an extraction standard, and lyophilized. Samples were then amended with 100 μ L of 0.01 M H_2SO_4 . After extraction, the

extracts were dried over anhydrous Na_2SO_4 , evaporated under a vacuum, and dissolved in 5 mL of ethyl acetate.

Parallel experiments utilizing [^{14}C]-dinoseb were performed to evaluate mass-balance relationships for dinoseb and its degradation products. Details are described below.

5 Results are shown in Table 3.

Table 3
Percent of Total Radioactivity

	Treatment	CO_2	Polar Extract	Nonpolar Extract	Solid	Percent Recovery
10	Loamy Sand Inoculated	29.6	43.3	5.11	4.0*	82
15	Loamy Sand Uninoculated	0.3	0.8*	100.8	6.7*	108
	Silt Loam Inoculated	32.1	29.9	3.5	8.9*	74
20	Silt Loam Uninoculated	0.8	1.1*	31.3	44.8	78

25 * = these values were not significantly different from background counts at the 90% confidence level

Referring to Table 3, about 30% mineralization to [^{14}C]- CO_2 was obtained after thirty days' incubation of samples of inoculated soils, as compared to less than 1% for the uninoculated controls. Most of the remaining radioactivity in the inoculated soils was present as polar metabolites. Less than 10% of the radioactivity was associated with solids in the inoculated soils after extraction, as compared to nearly 45% in the silt-loam control. In the loamy sand control, virtually all of the radioactivity was associated with the nonpolar extract, which contained undegraded dinoseb. In addition, although methane was not quantified in these experiments, some radioactivity appears to end up as methane.

35 To perform the radiochemical studies described above, radiolabeled dinoseb (u-ring [^{14}C]-dinoseb) was synthesized from [^{14}C]-phenol and was 96% radiochemically pure, as determined by HPLC analysis and TLC coupled with liquid scintillation counting. After adding 1 μCi u-ring [^{14}C]-dinoseb to the soil, the flasks were stoppered and a glass trap containing 1 mL of 1 M KOH was suspended in each flask. The KOH solution was exchanged daily during incubations, rinsed with 1 mL water and counted with 18 mL of BioSafe II liquid scintillation cocktail (RPI Inc., Mt. Prospect, IL). At the end of the incubation, each soil culture was connected to a CO_2 trap consisting of a series of four stoppered serum bottles each containing 10 mL of 1 N KOH. The gaseous effluent first passed through a C18 "Sep-Pak" cartridge (Waters Associates, Milford, MA) wetted with methanol to remove volatile organics. Each soil culture was then acidified with 10 mL concentrated HCl, agitated, and flushed with nitrogen gas to drive off dissolved CO_2 . One-mL samples from each trap were counted with 19 mL of BioSafe II.

Three 25-gram subsamples of each soil slurry were neutralized and extracted with 0.1 N NaOH. The extracts were neutralized, then passed through a "Sep-Pak" C18 cartridge which was rinsed with 1 mL water. The solid subsamples were dried, Soxhlet extracted, and the extracts were pooled with ethylacetate elutions from the "Sep-Pak" cartridges. From both the polar and non-polar extracts, 1 mL samples were counted with 19 mL of BioSafe II. Finally, 0.25 gm of each extracted soil sample was mixed with 19 mL BioSafe II and counted. All radioactive samples were counted in a Beckman Model 7000 liquid scintillation counter. A control flask for each soil type was treated similarly except that no bacterial inoculations were made.

In the above experiments, simply flooding the soils did not produce anaerobiosis in the resulting soil slurries, even in a slurry of the rich silt-loam soil. It is probable that anaerobiosis occurred within highly localized sites within the slurry, such as soil pores, but this was not sufficient to support growth of the oxygen-sensitive anaerobic consortium. In every case, exogenous carbon, preferably as a starchy carbohydrate metabolized by amylolytic microorganisms, was required to produce the requisite strict anaerobiosis that would allow growth of the subsequently added dinoseb-degrading anaerobic microorganisms.

The experiments described above show that exogenous, strictly anaerobic microorganisms can be used to degrade nitroaromatic chemicals in soil or wastewater. The mass balance and ^{14}C data indicate that the anaerobic consortium mediated a complete destruction of dinoseb as a representative nitroaromatic, rather than mere polymerization of the compound, which occurred in the silt-loam soil in the absence of anaerobic inoculation.

The method of the present invention, therefore, comprises basically two stages: an initial fermentative stage wherein the wastewater or aqueous slurry of nitroaromatic-contaminated soil is fortified with a starchy carbohydrate and inoculated with "fermentative" (aerobic and/or facultative) amylolytic bacteria that hydrolyze the starch, metabolize a portion of the sugars produced by the hydrolysis, and consume the oxygen in the liquid; and a subsequent anaerobic stage wherein the wastewater or aqueous slurry of contaminated soil is inoculated with a consortium of anaerobic bacteria that metabolize the remaining sugar and degrade the contaminant nitroaromatic. This bioremediation method may be used for any soil or water contaminated with dinoseb or other nitroaromatic or aminoaromatic compounds subject to polymerization reactions in aerobic soil, such as 4,6-dinitro-*o*-cresol and 3,5-dinitrobenzoate.

Although anaerobiosis can be performed in an open environment, it is inefficient and difficult to control unless performed in a closed environment, particularly on a large scale. As a result, the method of the present invention is performed in the field, preferably in a suitably large liquid-containment vessel or the like.

FIG. 14 is a schematic representation of one embodiment of the process of the present invention, as it would be conducted in the field to bioremediate a nitroaromatic-contaminated soil. In FIG. 14, the contaminated soil 22 is transferred via equipment 24 to a

-22-

plastic-lined pit 26 or the like. After the soil 22 is added to the pit 26, starchy carbohydrate 30, a nitrogen source 32, water 34, and an inoculum of fermentative amylolytic microorganism 36 are added to the soil 22, forming a slurry 28 in the pit 26. After forming the slurry 28, a cover 38 is placed over the pit 26, where the plastic-lined pit 26 and cover 38 together serve to exclude
5 gaseous exchange between the slurry 28 and the atmosphere. The slurry 28 is monitored using a potentiometric probe 40 coupled to a readout 42 to enable one to determine when anaerobic conditions have developed in the slurry 28 (redox potential about -200 mV or less). After the slurry 28 reaches anaerobiosis, an inoculum of an anaerobic consortium 44 of microorganisms is added to the slurry 28 beneath the surface, after performing sufficient numbers of serial transfers
10 46 of the anaerobic microorganisms 44 to yield a suitably large dose. The slurry 28 is maintained in an anaerobic condition in the covered pit until the nitroaromatics are biodegraded, at which time the slurry 28 may be returned 48 to the original site.

The process of FIG. 14 is a batch process. In a batch process applied to nitroaromatic-contaminated wastewater (not shown), the contaminated water is simply conducted
15 into the plastic-lined pit 26 of FIG. 14 or analogous vessel for bioremediation according to the present method. In the case of soil, it is usually necessary to add water to the soil to form a mud slurry, which is a more suitable environment for anaerobiosis than dry soil. Such a slurry will have a proportion of water relative to soil of about 15% to 20% (w/w) or more, depending upon the type of soil and the degree of nitroaromatic contamination, to preferably bring the soil at
20 least to about 100% water-holding capacity. More water can be added, if desired. More water added to the soil lessens the effective concentration of the contaminant nitroaromatic.

Since most soils and many natural waters already have a broad spectrum of mineral nutrients available to support life, in many cases it will not be necessary to add to the water or mud slurry many of the supplementary mineral nutrients required for laboratory cultures. In
25 virtually all cases, however, an extraneous nitrogen source 32 will be required, such as ammonium chloride. Many soils and most waters will also require extraneous phosphate (not shown), such as sodium or potassium phosphate. The nitrogen and phosphate requirements for the present process can be satisfied in many cases by merely adding ammonium phosphate (a common "fertilizer") as a supplement until the approximate desired concentration of ammonium
30 and phosphate are attained. Since each soil and water is different, preliminary analysis of the soil or water may be needed to determine indigenous concentrations of ammonium and phosphate so that the proper amount of supplementary ammonium and phosphate can be determined so as to avoid waste.

As stated above, the nitrogen source 32 should probably not be nitrate because
35 nitrates may inhibit the ability of the anaerobic microorganisms to biodegrade nitroaromatics.

It may be necessary, particularly in the case of strongly acidic waters or soils contaminated with nitroaromatics, to adjust the pH of the fluid medium to within the preferred range of about 6 to 8. An inexpensive additive, such as lime, is satisfactory for elevating the pH.

In most cases, however, significant pH adjustment will not be necessary. Extrapolating from the graph of FIG. 4, it would seem that pH values appreciably above or below (especially above) the preferred range of 6 to 8 would not cause a catastrophic depression of microbial metabolism required for bioremediation.

5 The preferred target concentrations of key nutrients (including contributions by the soil and/or water) are as given in the list of such ingredients in the nutrient solutions used in the studies described above. The ammonium concentration, as stated earlier, should preferably be between 0.4% and 0.5% (w/v), but a concentration between 0.1% and 0.6% (w/v) would suffice. The overall ionic strength of the water or slurry to be treated should preferably be approximately
10 that of the laboratory cultures of microorganisms, but the microbes are tolerant to surprisingly large variations of ionic strength.

 The temperature should preferably be between about 10°C and 40°C, with the optimal temperature about 25°C. As discussed above, lower non-freezing temperatures tend to slow microbial metabolism and correspondingly increase the length of time required to achieve
15 bioremediation.

 While yeast extract is an important supplement for laboratory cultures, it is not necessary in virtually all field applications, since soils and wastewater often contain "vitamins," cofactors, and other products of natural organic processes utilizable by the microorganisms of the present process.

20 The inoculum of fermentative amylolytic microorganisms 36 should be added at a dose of 10^7 to 10^8 CFU (colony forming units) per gram dry soil or per mL water. Higher doses are usually not necessary. Lower doses may result in a longer-than-optimal time to reach anaerobiosis. In addition, an amount of starchy carbohydrate 30, determined as described above, is also added. Preferably, the amount of starchy carbohydrate added per unit amount of water or
25 soil slurry is experimentally "tailored" to sustain anaerobiosis for a sufficient amount of time to achieve the desired degree of nitroaromatic degradation in the water or soil. Since types and concentrations of nitroaromatic contaminants will differ among various soils and waters, and since soils and waters themselves will differ, the optimal amount of starchy carbohydrate required will probably differ at each of various contaminated sites.

30 While the specific anaerobic consortium 44 of microorganisms described herein is suitable for degrading dinoseb and certain other nitroaromatics as described, other consortia of anaerobic microorganisms selected for and isolated in a manner as described herein, but using a nitroaromatic other than dinoseb may be more suitable for other nitroaromatics.

 Although employing fermentative microorganisms to render the water or slurry
35 anaerobic is the preferred method, other methods may be employed (not shown). However, it is anticipated that other methods may be prohibitively expensive. Such other methods include purging oxygen from the liquid using, for example, nitrogen or argon gas. However, gas-purging is typically slower and less efficient in achieving satisfactory anaerobic conditions than employing

aerobic fermentation. Another method would require adding oxygen-scavenging agents (strong reducing agents) to the liquid. Although use of reducing agents may be efficient (and rapid), the disadvantage is that such agents represent other contaminants added to the water or slurry. As a result, use of microbial fermentation of sugar or starch is the preferred approach for achieving anaerobiosis.

5 After the contaminant nitroaromatic has been satisfactorily biodegraded in the volume of water or slurry contained in the covered pit 26 or vessel, the pit 26 or vessel may be drained and a new volume of contaminated water or mud slurry added for bioremediation. In a semicontinuous process, about 10% to 15% of the previous batch of treated water or slurry may
10 be left in the pit 26 or vessel to aid in the inoculation of the subsequent batch. Additionally, such a semicontinuous process would preferably be controlled by various continuous electronic and chemical monitoring techniques known in the art, such as of dissolved oxygen and specific ions, as well as environmental conditions such as pH and temperature. Concentrations of nitroaromatic contaminants and their metabolic intermediaries can be discontinuously monitored using HPLC
15 and gas chromatography, for example.

As discussed above, the carbohydrate-fermentative microorganisms responsible for generation of anaerobic conditions and lowering of the redox potential of the fluid medium typically comprise various groups of such microorganisms present as a consortium. Likewise, the anaerobic microorganisms responsible for actual degradation of nitroaromatic compounds are
20 represented as a consortium of microorganisms. The species composition and species profiles of these consortia typically vary depending mainly upon the source of the microorganisms and upon the particular nitroaromatic compound to be degraded. (I.e., the inocula are preferably produced, as described above, by procedures in which microorganisms from a source are subjected to a "selection" protocol involving exposure to the particular nitroaromatic to be degraded. This
25 selection protocol as well as the inevitable variation in microbiological profiles of inocula obtained from different sources, result in the production of suitable inocula each represented by a distinctive species profile that functions as a consortium. Thus, various combinations of microorganisms can possess the complete metabolic capacity to degrade nitroaromatic compounds.

Despite the inherent variation in the species profile of various inocula, certain
30 groups of microorganisms routinely appear in each type of consortium. These groups are listed in Table 4, using nomenclature as found in Sneath et al. (eds.) Bergey's Manual of Systematic Bacteriology, Vol. 1-4, Williams and Wilkins, Baltimore (1986).

TABLE 4

I. Microorganisms typically present in consortia used for generation of anaerobic
35 conditions:

- A. Facultatively anaerobic Gram-negative rods of the family Enterobacteriaceae, including genera such as Klebsiella and Enterobacter.

-25-

- B. Gram-positive fermentative non-sporulating bacteria such as the genus Lactobacillus.
- C. Denitrifying bacteria such as the genera Bacillus, Clostridium, and Pseudomonas.

5

II. Microorganisms typically present in the anaerobic consortium:

- A. Microorganisms responsible for reduction of aromatic nitro groups to form reduced intermediates:
 - 1. Facultatively anaerobic Gram-negative rods of the family Enterobacteriaceae, including genera such as Klebsiella and Enterobacter.
 - 2. Anaerobic Gram-negative straight, curved, and helical rods of the family Bacteroidaceae, including genera such as Bacteroides and Fusobacterium.
 - 3. Dissimilatory sulfate- or sulfur-reducing bacteria, including genera such as Desulfovibrio and Desulfuromonas.
 - 4. Anaerobic endospore-forming Gram-positive rods and cocci, including genera such as Clostridium, Desulfotomaculum, and Sporosarcina.
 - 5. Gram-positive fermentative non-sporulating bacteria such as the genus Lactobacillus.
 - 6. Denitrifying bacteria, including genera such as Bacillus, Clostridium, and Pseudomonas.
- B. Microorganisms responsible for cleaving the aromatic rings of reduced intermediates of the nitroaromatics:
 - 1. Anaerobic Gram-negative straight, curved, and helical rods of the family Bacteroidaceae, including genera such as Bacteroides and Fusobacterium.
 - 2. Dissimilatory sulfate- or sulfur-reducing bacteria, including genera such as Desulfovibrio and Desulfuromonas.
 - 3. Anaerobic Gram-negative cocci, including genera such as Veillonella and Acidaminococcus.

10

15

20

25

30

35

-26-

4. Anaerobic endospore-forming Gram-positive rods and cocci, including genera such as Clostridium, Desulfotomaculum, and Sporosarcina.
- 5 C. Microorganisms responsible for anaerobic fermentation and anaerobic respiration and, ultimately, methanogenesis of the products of ring cleavage:
- 10 1. Anaerobic Gram-negative straight, curved, and helical rods of the family Bacteroidaceae, including genera such as Bacteroides and Fusobacterium.
2. Dissimilatory sulfate- or sulfur-reducing bacteria, including genera such as Desulfovibrio and Desulfuromonas.
- 15 3. Archaeobacteria, including Group-I methanogenic archaeobacteria; representative genera include Methanobacterium and Methanococcus.
- 20 4. Archaeobacteria, including Group-II archaeobacterial sulfate reducers; representative genus is Archaeoglobus.
5. Anaerobic endospore-forming Gram-positive rods and cocci, including genera such as Clostridium, Desulfotomaculum, and Sporosarcina.
- 25 6. Denitrifying bacteria, including genera such as Bacillus, Clostridium, and Pseudomonas.

30 Having illustrated and described the principles of our invention with reference to detailed descriptions of process steps and specific examples, it should be apparent to those of ordinary skill in the art that the invention may be modified in arrangement and detail without departing from such principles. We claim as our invention all such modifications as come within the true spirit and scope of the following claims.

Claims

1. A method for degrading nitroaromatic compounds present as contaminants in a fluid medium to reduce the concentrations of the nitroaromatic compounds in the fluid medium, the method comprising:

5 (a) adding to the fluid medium an inoculum of carbohydrate-fermentative microorganisms resistant to the types and respective concentrations of contaminant nitroaromatic compounds present in the fluid medium, wherein the carbohydrate-fermentative microorganisms are selected from the group consisting of aerobic and facultative microorganisms comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Lactobacillus*, *Bacillus*,
10 *Clostridium*, and *Pseudomonas* and mixtures thereof;

(b) adding to the fluid medium a carbohydrate fermentable by the carbohydrate-fermentative microorganisms;

(c) maintaining the fluid medium containing said carbohydrate-fermentative microorganisms and carbohydrate under conditions suitable for growth of said carbohydrate-fermentative microorganisms in the fluid medium, accompanied by fermentation by said
15 microorganisms of a portion of said carbohydrate, thereby depleting the concentration of dissolved oxygen in the fluid medium, so as to render the fluid medium anaerobic;

(d) after the fluid medium has been rendered anaerobic, adding to the fluid medium an inoculum of anaerobic microorganisms capable of utilizing the carbohydrate as a carbon and energy source and degrading said nitroaromatic compounds, the anaerobic inoculum being a consortium comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Bacteroides*, *Fusobacterium*, *Desulfovibrio*, *Desulfuromonas*, *Clostridium*, *Desulfotomaculum*, *Sporosarcina*, *Lactobacillus*, *Bacillus*, *Clostridium*, *Pseudomonas*, *Veillonella*, *Acidaminococcus*, *Methanobacterium*, *Methanococcus*, and *Archaeoglobus* and mixtures thereof; and
20

(e) maintaining anaerobic conditions in the fluid medium for a time sufficient for the anaerobic microorganisms to degrade the contaminant nitroaromatic compounds.
25

2. A method for degrading nitroaromatic compounds as recited in claim 1, including the step of adding mineral nutrients to the fluid medium to support growth of the microorganisms inoculated into the fluid medium.

3. A method for degrading nitroaromatic compounds as recited in claim 1, including the step of adding a source of metabolizable nitrogen to the fluid medium to support the growth of the microorganisms inoculated into the fluid medium.
30

4. A method for degrading nitroaromatic compounds as recited in claim 3, wherein the source of metabolizable nitrogen is ammonium chloride.

35 5. A method for degrading nitroaromatic compounds as recited in claim 4, wherein the ammonium chloride is added to the fluid medium to yield a concentration of ammonium chloride in the fluid medium within a range of about 4 g/L to 5 g/L.

6. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the fluid medium is maintained at a pH within a range of about 6 to 8.

7. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the fluid medium is maintained at a temperature within a range of about 10°C to less than about 40°C.

8. A method for degrading nitroaromatic compounds as recited in claim 1, wherein said anaerobic microorganisms are capable of degrading dinoseb present as a contaminant in the fluid medium.

9. A method for degrading nitroaromatic compounds as recited in claim 8, wherein the inoculum of carbohydrate-fermentative microorganisms includes *Klebsiella oxytoca* strain DSA-1.

10. A method for degrading nitroaromatic compounds as recited in claim 9, wherein said *Klebsiella oxytoca* strain DSA-1 microorganisms are resistant to dinoseb at a concentration of up to at least 100 ppm.

11. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the carbohydrate-fermentative microorganisms are added at an inoculum dosage within a range of at least about 10^7 to 10^8 CFU per gram of fluid medium.

12. A method for degrading nitroaromatic compounds as recited in claim 1, wherein said anaerobic microorganisms are capable of degrading nitrotoluene compounds present as contaminants in the fluid medium.

13. A method for degrading nitroaromatic compounds as recited in claim 12, wherein said nitrotoluene compounds present as contaminants in the fluid medium include trinitrotoluene.

14. A method for degrading nitroaromatic compounds as recited in claim 1, wherein formation of polymerized derivatives of said contaminant nitroaromatic compounds in the fluid medium is inhibited.

15. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the inoculum of carbohydrate-fermentative microorganisms is comprised of a population of microorganisms grown in aerobic nutrient medium including the nitroaromatic compounds present as contaminants in said fluid medium.

16. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the redox potential of the fluid medium when anaerobic is less than about -200 mV.

17. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the carbohydrate is comprised of a starch.

18. A method for degrading nitroaromatic compounds as recited in claim 17, wherein said carbohydrate is comprised of a dewatered potato by-product.

19. A method for degrading nitroaromatic compounds as recited in claim 17, wherein the carbohydrate-fermentative microorganisms are amylolytic.

20. A method for degrading nitroaromatic compounds as recited in claim 1, wherein the inoculum of anaerobic microorganisms is comprised of a population of anaerobic microorganisms grown in nutrient medium including the nitroaromatic compounds present as contaminants in said fluid medium.

5 21. A method for degrading nitroaromatic compounds as recited in claim 1, wherein at least a portion of said nitroaromatic compounds is degraded to carbon dioxide.

22. A method for degrading nitroaromatic compounds as recited in claim 1, including the steps of placing the nitroaromatic-contaminated fluid medium in a covered vessel before rendering the fluid medium anaerobic and keeping the fluid medium contained within the covered vessel until said anaerobic microorganisms have been added to the vessel and have degraded the nitroaromatic compounds in the fluid medium.

23. A method for remediating a soil so as to degrade nitroaromatic compounds present as contaminants in the soil, the method comprising:

15 (a) adding water to the nitroaromatic-contaminated soil to form an aqueous slurry of the soil;

(b) adding to the slurry an inoculum of fermentative amylolytic microorganisms resistant to the nitroaromatic compounds and concentrations thereof present in the soil, wherein the fermentative amylolytic microorganisms are selected from the group consisting of aerobic and facultative amylolytic microorganisms comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Lactobacillus*, *Bacillus*, *Clostridium*, and *Pseudomonas* and mixtures thereof;

(c) adding to the slurry at least one type of starchy carbohydrate hydrolyzable to substituent sugars by said fermentative amylolytic microorganisms;

25 (d) maintaining the slurry containing said fermentative amylolytic microorganisms and starchy carbohydrate under conditions suitable for growth of said fermentative amylolytic microorganisms in the slurry, accompanied by hydrolysis of the starchy carbohydrate to constituent sugars and metabolism by said fermentative amylolytic microorganisms of a portion of said sugars, thereby rendering the slurry anaerobic;

30 (e) after the slurry has been rendered anaerobic, adding to the slurry an inoculum of anaerobic microorganisms capable of utilizing as a carbon and energy source said sugars formed in the slurry by the amylolytic microorganisms and capable of degrading said nitroaromatic compounds the anaerobic inoculum being a consortium comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Bacteroides*, *Fusobacterium*, *Desulfovibrio*, *Desulfuromonas*, *Clostridium*, *Desulfotomaculum*, *Sporosarcina*, *Lactobacillus*, *Bacillus*, *Clostridium*, *Pseudomonas*, *Veillonella*, *Acidaminococcus*, *Methanobacterium*, *Methanococcus*, and *Archaeoglobus* and mixtures thereof; and

(f) maintaining anaerobic conditions in the slurry for a time sufficient for the anaerobic microorganisms to degrade the nitroaromatic compounds.

24. A method for remediating soil as recited in claim 23, wherein the water is added to the soil in a volume sufficient to saturate the water-holding capacity of the soil.

25. A method for remediating soil as recited in claim 23, including the step of adding a source of metabolizable nitrogen to the slurry to support the growth of the
5 microorganisms inoculated into the slurry.

26. A method for remediating soil as recited in claim 24, wherein the source of metabolizable nitrogen is ammonium chloride.

27. A method for remediating soil as recited in claim 25, wherein the ammonium chloride is added to a concentration in the water within a range of about 4 g/L to about 5 g/L.

10 28. A method for remediating soil as recited in claim 23, wherein the slurry is maintained at a pH within a range of about 6 to 8.

29. A method for remediating soil as recited in claim 23, wherein the slurry is maintained at a temperature within a range of about 10°C to less than about 40°C.

15 30. A method for remediating soil as recited in claim 23, wherein the inoculum of fermentative amylolytic microorganisms is comprised of a population of microorganisms grown in aerobic nutrient medium including the nitroaromatic compounds present as contaminants in said soil.

31. A method for remediating soil as recited in claim 23, wherein the redox potential of the slurry when anaerobic is less than about -200 mV.

20 32. A method for remediating soil as recited in claim 23, wherein the inoculum of anaerobic microorganisms is comprised of a population of anaerobic microorganisms grown in nutrient medium, including the nitroaromatic compounds present as contaminants in said soil.

33. A method for remediating soil as recited in claim 23, wherein at least a portion of said contaminant nitroaromatic compounds is degraded to carbon dioxide.

25 34. A method for remediating soil as recited in claim 23, including the steps of placing the nitroaromatic-contaminated soil in a vessel before adding said water to the soil to form said slurry, covering the vessel containing the slurry so as to substantially exclude gaseous exchange between the slurry and the atmosphere, and keeping the slurry contained within the covered vessel until said anaerobic microorganisms have degraded the nitroaromatic compounds
30 in the slurry.

35. A method for treating a soil contaminated with aminoaromatic compounds so as to favor mineralization of the aminoaromatic compounds in the soil over oxidative polymerization of the aminoaromatic compounds, the method comprising:

35 (a) adding water to the nitroaromatic-contaminated soil to form an aqueous slurry of the soil;

(b) adding to the slurry an inoculum of fermentative amylolytic microorganisms resistant to the aminoaromatic compounds in the soil, where the fermentative amylolytic microorganisms are selected from the group consisting of aerobic and facultative amylolytic

microorganisms comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Lactobacillus*, *Bacillus*, *Clostridium*, and *Pseudomonas* and mixtures thereof;

(c) adding to the slurry at least one type of starchy carbohydrate hydrolyzable to constituent sugars by said fermentative amylolytic microorganisms;

5 (d) maintaining the slurry containing said inoculum of fermentative amylolytic microorganisms and starchy carbohydrate under conditions suitable for growth of said fermentative microorganisms in the slurry, accompanied by hydrolysis of the starchy carbohydrate to constituent sugars and metabolism by said microorganisms of a portion of said sugars, thereby depleting the concentration of dissolved oxygen in the slurry so as to render the slurry anaerobic;

10 (e) after the slurry has been rendered anaerobic, adding to the slurry an inoculum of anaerobic microorganisms capable of utilizing as a carbon and energy source said sugars formed in the slurry by the amylolytic microorganisms and capable of degrading said aminoaromatic compounds, the anaerobic inoculum being a consortium comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Bacteroides*, *Fusobacterium*,
15 *Desulfovibrio*, *Desulfuromonas*, *Clostridium*, *Desulfotomaculum*, *Sporosarcina*, *Lactobacillus*, *Bacillus*, *Clostridium*, *Pseudomonas*, *Veillonella*, *Acidaminococcus*, *Methanobacterium*, *Methanococcus*, and *Archaeoglobus* and mixtures thereof; and

(f) maintaining anaerobic conditions in the slurry for a time sufficient for the anaerobic microorganisms to degrade the contaminant aminoaromatic compounds.

20 36. A method for treating soil as recited in claim 35, wherein the water is added to the soil in a volume sufficient to saturate the water-holding capacity of the soil.

37. A method for treating soil as recited in claim 35, including the step of adding a source of metabolizable nitrogen to the slurry to support the growth of the microorganisms inoculated into the slurry.

25 38. A method for treating soil as recited in claim 35, wherein the slurry is maintained at a pH within a range of about 6 to 8.

39. A method for treating soil as recited in claim 35, wherein the slurry is maintained at a temperature within a range of about 10°C to less than about 40°C.

30 40. A method for treating soil as recited in claim 35, wherein the redox potential of the slurry when anaerobic is less than about -200 mV.

41. A method for treating soil as recited in claim 35, wherein the inoculum of fermentative amylolytic microorganisms is comprised of a population of amylolytic microorganisms grown in aerobic nutrient medium including the aminoaromatic compounds present as contaminants in said soil.

35 42. A method for treating soil as recited in claim 35, wherein the inoculum of anaerobic microorganisms is comprised of a population of anaerobic microorganisms grown in nutrient medium including the aminoaromatic compounds present as contaminants in said soil.

43. A method for treating soil as recited in claim 35, including the steps of placing the aminoaromatic-contaminated soil in a vessel before adding said water to the soil to form said slurry, covering the vessel containing the slurry so as to substantially exclude gaseous exchange between the slurry and the atmosphere, and keeping the slurry contained in the covered vessel until said anaerobic microorganisms have degraded the aminoaromatic compounds in the slurry.

44. A two-stage process for treating a fluid medium contaminated with nitroaromatic compounds so as to at least partially degrade and thereby reduce the concentrations of the nitroaromatic compounds in the fluid medium, the process comprising:

(a) a fermentative stage wherein a first inoculum is added to the fluid medium, wherein the first inoculum comprises (1) carbohydrate-fermentative microorganisms resistant to the concentration of nitroaromatic compounds present in the fluid medium, and wherein the carbohydrate-fermentative microorganisms are selected from the group consisting of aerobic and facultative microorganisms comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Lactobacillus*, *Bacillus*, *Clostridium*, and *Pseudomonas* and mixtures thereof, (2) at least one type of carbohydrate fermentable by said carbohydrate-fermentative microorganisms, and (3) a source of metabolizable nitrogen, after which addition the fluid medium is maintained under conditions suitable for growth of the carbohydrate-fermentative microorganisms in the fluid medium, accompanied by fermentation by said microorganisms of a portion of said carbohydrate, thereby depleting the concentration of dissolved oxygen in the fluid medium so as to render the fluid medium anaerobic; and

(b) an anaerobic stage after the fermentative stage, in which anaerobic stage is added to the fluid medium a second inoculum, wherein the second inoculum comprises anaerobic microorganisms capable of utilizing said carbohydrate remaining from the fermentative stage as a carbon and energy source and of degrading said nitroaromatic compounds, the second inoculum being a consortium comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Bacteroides*, *Fusobacterium*, *Desulfovibrio*, *Desulfuromonas*, *Clostridium*, *Desulfotomaculum*, *Sporosarcina*, *Lactobacillus*, *Bacillus*, *Clostridium*, *Pseudomonas*, *Veillonella*, *Acidaminococcus*, *Methanobacterium*, *Methanococcus*, and *Archaeoglobus* and mixtures thereof, after which addition the fluid medium is maintained in an anaerobic condition for a time sufficient for the anaerobic microorganisms to degrade the contaminant nitroaromatic compounds in the fluid medium.

45. A two-stage process as recited in claim 44, wherein the fluid medium is maintained at a pH within a range of about 6 to 8.

46. A two-stage process as recited in claim 44, wherein the fluid medium is maintained at a temperature within a range of about 10°C to less than about 40°C.

47. A two-stage process as recited in claim 44, wherein the source of metabolizable nitrogen is selected from the group consisting of ammonium compounds and amino compounds.

48. A two-stage process as recited in claim 44, wherein the first inoculum is comprised of a population of microorganisms grown in aerobic nutrient medium including the nitroaromatic compounds present as contaminants in said fluid medium.

5 49. A two-stage process as recited in claim 44, wherein the carbohydrate comprises a starch.

50. A two-stage process as recited in claim 49, wherein the first inoculum includes amylolytic microorganisms.

10 51. A two-stage process as recited in claim 44, wherein the second inoculum is comprised of a population of anaerobic microorganisms grown in nutrient medium including the nitroaromatic compounds present as contaminants in said fluid medium.

52. A method for degrading nitroaromatic compounds present as contaminants in a fluid medium to reduce the concentrations of the nitroaromatic compounds in the fluid medium, the method comprising:

15 (a) sequestering the fluid medium from the atmosphere so as to prevent gaseous exchange between the fluid medium and the atmosphere;

(b) rendering the fluid medium anaerobic;

20 (c) adding to the fluid medium an inoculum of anaerobic microorganisms capable of degrading the nitroaromatic compounds in the fluid medium the anaerobic inoculum being a consortium comprising genera selected from the group consisting of *Klebsiella*, *Enterobacter*, *Bacteroides*, *Fusobacterium*, *Desulfovibrio*, *Desulfuromonas*, *Clostridium*, *Desulfotomaculum*, *Sporosarcina*, *Lactobacillus*, *Bacillus*, *Clostridium*, *Pseudomonas*, *Veillonella*, *Acidaminococcus*, *Methanobacterium*, *Methanococcus*, and *Archaeoglobus* and mixtures thereof;

(d) adding to the fluid medium metabolizable sources of carbon, energy, and nitrogen for the anaerobic microorganisms inoculated into the fluid medium;

25 (e) maintaining the fluid medium in an anaerobic condition for a time sufficient for the anaerobic microorganisms to degrade the contaminant nitroaromatic compounds in the fluid medium.

53. A method for degrading nitroaromatic compounds in a fluid medium as recited in claim 52, wherein the fluid medium is rendered anaerobic by purging oxygen from the fluid medium.

30 54. A method for degrading nitroaromatic compounds in a fluid medium as recited in claim 52, wherein the fluid medium is rendered anaerobic by adding to the fluid medium an oxygen-scavenging agent.

35 55. A method for degrading nitroaromatic compounds in a fluid medium as recited in claim 52, wherein the fluid medium is sequestered from the atmosphere by placing the fluid medium in a vessel for containing the fluid medium and covering the vessel.




FIG. 1

2 / 10




FIG. 2




FIG. 3




FIG. 4

4 / 10




FIG. 5




FIG. 6

5 / 10




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11

8 / 10




FIG. 12

9 / 10





FIG. 13

FIG. 14



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/08431

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : C05F 11/08; C02F 03/00; C05G 03/00

US CL : 71/6,9,10,903; 435/262,262.5; 210/610,611

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 71/6,9,10,903,904; 435/262,262.5; 210/610,611

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US,A, 4,925,552 (BATESON ET AL) 15 MAY 1990, SEE ENTIRE DOCUMENT	1-55

☐ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

Special categories of cited documents:	
A document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
E earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
O document referring to an oral disclosure, use, exhibition or other means	*Z* document member of the same patent family
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

26 OCTOBER 1994

Date of mailing of the international search report

NOV 07 1994

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized Officer

PERRIS H. LANDER

Telephone No. (703) 308-0533

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☒ FADED TEXT OR DRAWING
- ☒ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☐ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.